



## Fish energy budget under ocean warming and flame retardant exposure

Patrícia Anacleto<sup>a,b,c,\*</sup>, Cátia Figueiredo<sup>b</sup>, Miguel Baptista<sup>b</sup>, Ana Luísa Maulvault<sup>a,b,c</sup>,  
 Carolina Camacho<sup>a,c,d</sup>, Pedro Pousão-Ferreira<sup>a</sup>, Luísa M.P. Valente<sup>c,e</sup>, António Marques<sup>a,c</sup>,  
 Rui Rosa<sup>b</sup>

<sup>a</sup> Division of Aquaculture and Upgrading (DivAV), Portuguese Institute for the Sea and Atmosphere (IPMA, I.P.), Rua Alfredo Magalhães Ramalho 6, 1495-006 Lisboa, Portugal

<sup>b</sup> MARE – Marine and Environmental Sciences Centre, Guia Marine Laboratory, Faculty of Sciences, University of Lisbon (FCUL), Av. Nossa Senhora do Cabo, 939, 2750-374 Cascais, Portugal

<sup>c</sup> Interdisciplinary Centre of Marine and Environmental Research (CIIMAR), University of Porto, Terminal de Cruzeiros do Porto de Leixões, Av. General Norton de Matos, S/N, 4450-208 Matosinhos, Portugal

<sup>d</sup> Faculty of Sciences and Technology, New University of Lisbon, Quinta da Torre, 2829-516 Caparica, Portugal

<sup>e</sup> ICBAS, Instituto de Ciências Biomédicas de Abel Salazar, University of Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal



### ARTICLE INFO

#### Keywords:

Ocean warming  
 BDE-209  
 White seabream  
 Energy budget  
 Metabolism

### ABSTRACT

Climate change and chemical contamination are global environmental threats of growing concern for the scientific community and regulatory authorities. Yet, the impacts and interactions of both stressors (particularly ocean warming and emerging chemical contaminants) on physiological responses of marine organisms remain unclear and still require further understanding. Within this context, the main goal of this study was to assess, for the first time, the effects of warming (+ 5 °C) and accumulation of a polybrominated diphenyl ether congener (BDE-209, brominated flame retardant) through dietary exposure on energy budget of the juvenile white seabream (*Diplodus sargus*). Specifically, growth (G), routine metabolism (R), excretion (faecal, F and nitrogenous losses, U) and food consumption (C) were calculated to obtain the energy budget. The results demonstrated that the energy proportion spent for G dominated the mode of the energy allocation of juvenile white seabream (56.0–67.8%), especially under the combined effect of warming plus BDE-209 exposure. Under all treatments, the energy channelled for R varied around 26% and a much smaller percentage was channelled for excretion (F: 4.3–16.0% and U: 2.3–3.3%). An opposite trend to G was observed to F, where the highest percentage (16.0 ± 0.9%) was found under control temperature and BDE-209 exposure via diet. In general, the parameters were significantly affected by increased temperature and flame retardant exposure, where higher levels occurred for: i) wet weight, relative growth rate, protein and ash contents under warming conditions, ii) only for O:N ratio under BDE-209 exposure via diet, and iii) for feed efficiency, ammonia excretion rate, routine metabolic rate and assimilation efficiency under the combination of both stressors. On the other hand, decreased viscerosomatic index was observed under warming and lower fat content was observed under the combined effect of both stressors. Overall, under future warming and chemical contamination conditions, fish energy budget was greatly affected, which may dictate negative cascading impacts at population and community levels. Further research combining other climate change stressors (e.g. acidification and hypoxia) and emerging chemical contaminants are needed to better understand and forecast such biological effects in a changing ocean.

### 1. Introduction

Climate change is one of the greatest environmental challenges that cause profound and diverse impacts on marine organisms and ecosystems (IPCC, 2014). Anthropogenic activities are strongly influencing climate, primarily through fossil fuels, industrial, agricultural, and other land use emissions that alter atmospheric carbon dioxide (CO<sub>2</sub>).

Hence, one direct consequence of cumulative and continuing emissions of CO<sub>2</sub> since the industrial revolution is the increase of surface ocean temperatures (Doney et al., 2012). According to the latest report of the Intergovernmental Panel on Climate Change (IPCC), the average temperatures on the Earth's surface are expected to rise between 0.3 °C and 4.8 °C by the end of the 21st century with strong local variations. Additionally, it is predicted that extreme high temperature events (heat

\* Correspondence to: Portuguese Institute of Sea and Atmosphere (IPMA, I.P.), Rua Alfredo Magalhães Ramalho 6, 1495-006 Lisboa, Portugal.  
 E-mail address: [panacleto@ipma.pt](mailto:panacleto@ipma.pt) (P. Anacleto).

waves) will become more extreme and frequent (IPCC, 2014). Ocean warming represents a major threat to many marine organisms deeply affecting their physiological responses, namely growth, feeding, acid–base balance, metabolism and behaviour in ways that often compromise species fitness and survival (Pörtner and Peck, 2010; Rosa et al., 2012a). Thus, temperature is one of the main abiotic factors that can lead to significant changes on the energy budget of an organism (Yurista, 1999). The energy budget describes the energy intake and expenditure within the whole organism, leading to the partitioning of ingested energy into the major physiological components of the energy budget equation:  $C = G + R + F + U$ , where energy consumed in form of food (C) is the sum of energy retained in materials deposited during growth (G), energy spent for respiration (R), energy losses through faeces (F) and energy losses via nitrogenous excretory products (U; non-faecal) (Jobling, 1994). At optimal temperature conditions, an organism allocates a great part of its energy input into vital functions, as well as in the build-up of energy storage. In contrast, under stress (sub-optimal high temperatures), fish cannot consume enough food to meet, for example, the increasing metabolic costs, whereby energy is often limited and can only cover costs of essential functions, such as standard metabolism (Clarke and Johnston, 1999). Indeed, warming greatly affect metabolic rate of fish, since there is a dependent relation between the variables (temperature–oxygen) and oxygen supply becomes limited at high temperature borders (Pörtner and Peck, 2010; Rosa et al., 2014). Consequently, all of biological processes are directly affected by temperature, namely food intake, nutritional efficiency and whole-body composition (Burel et al., 1996; Fang et al., 2010). Zheng et al. (2008) also reported that growth rate and faecal production increased in fish species at higher temperatures, while nitrogen excretion decreased, leading to a higher O:N ratio (ratio of oxygen consumed to nitrogen excreted). On the other hand, ocean warming impacts are also expected to directly influence the availability and toxicological effects of marine pollutants (Marques et al., 2010). Recent studies, evidenced that marine fish species tend to accumulate emerging chemical contaminants in warmer conditions (Maulvault et al., 2016, 2017). Such trend was demonstrated in our recent work with seabass (*Dicentrarchus labrax*) juveniles exposed to methylmercury (MeHg) via diet, with specimens exposed to warmer temperatures (22 °C) showing higher levels of MeHg in different tissues (muscle, liver and brain; Maulvault et al., 2016). The contaminants of emerging concern are a subject of growing interest for the scientific community and regulatory authorities, since constitute a risk for human health and biota, and for which there is still insufficient scientific knowledge (Cruz et al., 2015). Among these contaminants, polybrominated diphenyl ethers (PBDEs) are an important class of flame retardants (FRs) consisting of 209 congeners, which have been commercialized as penta-, octa-, and decabrominated mixtures. PBDEs are additive flame retardants, i.e. they are not bound to the polymer, but have been widely used in a variety of products (e.g. plastics, furniture, vehicles) to make them less combustible and to retard flames (Alaee et al., 2003). These FRs are persistent in the environment, accumulate in food chains and show toxic effects on hormonal regulation and affect neuronal, thyroid and liver-related activities (Costa and Giordano, 2011). Hence, these substances are the subject of a wide range of toxicologic and ecotoxicologic studies. Due to these concerns of environmental persistence, bioaccumulation and toxicity, lower brominated PBDEs have been banned in several countries, but the congener decabromodiphenyl ether (BDE-209), primary constituent of deca-BDE, is still used. Nonetheless, it is worth noting that some European countries (e.g. Norway) or states in the U.S.A. (e.g. Maine) have recently banned their use (Costa and Giordano, 2011). Despite the lower levels of BDE-209 found in water, seafood has been identified as the major dietary source of this compound for human uptake, since it can be biotransformed to lower and more toxic congeners via the food web (Kierkegaard et al., 1999; Elliott et al., 2005; Stapleton et al., 2006). Several studies have addressed the effects of water temperature on energy budget of some fish species (e.g. Xie et al., 2011), but the

conclusions were not consistent among them, and others only focused on the effects of chemical contaminants under climate change (e.g. Maulvault et al., 2016, 2017; Sampaio et al., 2018) without ever reporting the effects on energy budgets. Yet, a small number of studies investigated the effects of BDE-209 dietary exposure only on growth of marine species (Kuo et al., 2010; Zhang et al., 2013; Sha et al., 2015) and, to our knowledge, there are no studies that investigated the interactive effects of both stressors (ocean warming and the presence of emerging chemical contaminants) in the perspective of fish bioenergetics. In this context, the main goal of this study was to investigate, for the first time, the effects of ocean warming (+ 5 °C) and the dietary exposure to the emerging BDE-209 (60 ng g<sup>-1</sup> dry weight, dw) on the energy budget of juvenile white seabream (*Diplodus sargus*), by measuring growth, routine metabolism, excretion and food consumption. This species was selected as a suitable biological model to evaluate these impacts since it generally inhabits coastal areas particularly susceptible to hydrographic changes and where climate change effects may certainly pose greater ecological and toxicological challenges (Marques et al., 2010; Rosa et al., 2012b). Additionally, this seabream species plays an important role in the coastal food web, as they link top predators with species occupying low trophic levels.

## 2. Materials and methods

### 2.1. Experimental design and sampling

Juvenile white seabream (*Diplodus sargus*) specimens from the same batch and with similar biometric characteristics (6.2 ± 0.6 cm total length; 3.9 ± 1.2 g total weight; mean ± standard deviation) were reared at the aquaculture pilot station of the Portuguese Institute for the Sea and Atmosphere (EPPO-IPMA, Olhão, Portugal) and transported in thermal isolated containers, with constant aeration, to Guia Marine Laboratory (MARE-FCUL, Cascais, Portugal). Upon arrival, fish were randomly distributed in 12 rectangular incubating glass tanks (98 × 33 × 24.7 cm; 100 L total capacity each), each with independent recirculating aquaculture systems (RAS). Each RAS was equipped with biological filtration (model FSBF 1500, TMC Iberia, Portugal), physical filtration (protein skimmer; ReefSkinPro, TMC-Iberia, Portugal), UV disinfection (Vecton 300, TMC Iberia, Portugal), independent and automatic temperature (Fimar, Fernando Ribeiro Lda, Portugal) and pH control (model Profilux 3.1 N, GHL, Germany) via solenoid valves system. Each incubation system (n = 12) had independent pH measurements, by means of pH electrodes (n = 24 in total), connected to the Profilux system apparatus. pH values were monitored every 2 s and lowered by injection of a certified CO<sub>2</sub> gas mixture (Air Liquide, Portugal) via air stones or increased by tank aeration with CO<sub>2</sub>-filtered (using soda lime, Sigma-Aldrich) air. Seawater used in the RAS was filtered (0.35 µm) and UV sterilized (Vecton 600, TMC Iberia, Portugal). Ammonia (NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup>), nitrite (NO<sub>2</sub><sup>-</sup>) and nitrate (NO<sub>3</sub><sup>-</sup>) levels were monitored regularly, by means of colorimetric tests kits (Tropic Marin, USA) and kept below 0.05 mg L<sup>-1</sup>, 0.20 mg L<sup>-1</sup> and 2.0 mg L<sup>-1</sup>, respectively. In order to avoid physiological stress related to high animal density, fish density was kept below 5 g body weight L<sup>-1</sup> in each incubation tank. Daily, mortality was registered in each treatment and seawater was partially replaced (around 20%).

Fish were acclimated to laboratory conditions during two weeks before the start of the experiment at the following abiotic conditions: dissolved oxygen (DO) above 5 mg L<sup>-1</sup>; temperature = 19 ± 0.4 °C; pH = 8.06 ± 0.10; salinity = 35 ± 1‰ (WTW handheld Meter Multi 350i, Germany) and photoperiod of 12 h light and 12 h dark (12 L:12D). Five days before initiating the climate change exposure scenarios, seawater temperature was slowly raised (1 °C per day) until reaching 24 °C in tanks simulating warming conditions, in order to allow specimens to acclimate at this temperature before the beginning of the trial. No mortality occurred during the acclimation period.

Fish were exposed to four scenarios during 56 days to understand

future ocean changes expected for 2100 (i.e. seawater warming,  $\Delta = 5^\circ\text{C}$  according to IPCC projections scenario RCP8.5; IPCC, 2014): i) Scenario 1 (S1) –  $19^\circ\text{C}_{\text{Control}}$ , i.e. seawater temperature set at  $19^\circ\text{C}$  (control; current conditions used in juvenile seabream rearing in Iberian Peninsula) and animals fed with the control diet; ii) Scenario 2 (S2) –  $24^\circ\text{C}_{\text{Control}}$  – seawater temperature set at  $24^\circ\text{C}$  (warming scenario) and animals fed with the control diet; iii) Scenario 3 (S3) –  $19^\circ\text{C}_{\text{BDE-209}}$  exposure – seawater temperature set at  $19^\circ\text{C}$  and animals fed with BDE-209 contaminated diet; and iv) Scenario 4 (S4) –  $24^\circ\text{C}_{\text{BDE-209}}$  exposure – seawater temperature set at  $24^\circ\text{C}$  (warming scenario) and animals fed with BDE-209 contaminated diet. Each treatment comprised three independent replicate tanks ( $n = 17$  animals per replicate/tank, i.e. 51 animals per treatment). During the experimental period, specimens were fed with 2% of the average body weight (divided in two meals per day) with the respective diets (non-contaminated for control treatments and during acclimation period for all treatments, and contaminated with BDE-209 for contaminated treatments during the trial). Individual weekly food rations were adjusted according to fish size and biomass in each tank throughout the experiment. Before each sampling, fish were fasted for 24 h and anaesthetized using tricaine methanesulfonate ( $70\text{ mg L}^{-1}$  of MS-222; Sigma-Aldrich, St Louis, MO, USA) buffered with sodium bicarbonate ( $\text{NaHCO}_3$ ; Sigma-Aldrich) using a ratio of 1:1 to reduce fish stress.

In the beginning and end of the experiment (T0 and T56), all fish were fasted for 24 h to allow evacuation of faeces before bulk-weighing. Additionally, fish were weighed weekly to adjust feed rations in all treatments. Fish trials were approved by the Ethical Committee of the Faculty of Sciences of the Lisbon University (ORBEA) and conducted according to legal regulations (EU Directive 2010/63). Five specimens per replicate/tank ( $n = 15$  per treatment) were randomly weighted and sampled at the beginning and end of the experiment and frozen at  $-20^\circ\text{C}$  for chemical analysis and assessment of energy content. All samples (whole fish) were freeze-dried for 48 h, at  $-50^\circ\text{C}$  with low vacuum pressure (approximately  $10^{-1}$  atm; Power Dry 150 LL3000, Heto, Czech Republic), homogenized and kept at  $-80^\circ\text{C}$  until further analyses. Also, biometric data (total weight and total length) was registered at T0 and T56 (Table 2).

## 2.2. Feed

Non-contaminated (control) and contaminated dry inert pellets (BDE-209-spiked) with the same nutritional composition were manufactured by a specialized feed producing company (SPAROS Lda, Olhão, Portugal), considering the nutritional requirements for juvenile white seabream (detailed formulation and chemical composition can be consulted in Table A.1). Briefly, all ingredients were grinded (below  $200\ \mu\text{m}$ ) in a micropulverizer hammer mill (Hosokawa Micron, SH1, The Netherlands). Ingredients and fish oil were then mixed accordingly to the target formulation in a paddle mixer (Mainca RM90, Spain), and further humidified with 25% deionized water at room temperature. The diet was extruded at  $3.0\text{ mm}$  by means of a low-shear extruder (P55, Italplast, Italy). Upon extrusion, feed pellets were dried in a vibrating fluid bed dryer (model DR100, TGC Extrusion, France). A batch of this control diet was contaminated with BDE-209. A contaminant stock solution was prepared beforehand, by solubilising the contaminant in a selenium solution ( $250\ \mu\text{g/mL}$ ) and further diluted with acetone, in order to achieve a BDE-209-enriched feed with a nominal concentration of  $60\text{ ng g}^{-1}$  on a dry weight basis (dw). This stock solution was further mixed in fish oil (total volume of  $200\text{ mL}$ ) and this solution was top-coated to the pellets with a pressurized spraying container (standard flat-fan nozzle; size  $250\ \mu\text{m}$ ; pressure  $3.1\text{ bar}$ ). Contaminated diet was stored in a dark and cool place during its usage (about 1.5 month). Contamination level of this PBDE congener was systematically assessed during the experiment, by quantification of BDE-209 content in both feeds, using a gas chromatography-mass spectrometry (GC-MS; Agilent 7890B GC coupled to an Agilent 7000 C triple quadrupole mass

spectrometer), as described in detail by Cruz et al. (2017). The concentration of BDE-209 in control and contaminated diets were below limit of detection (LOD) and  $60.52 \pm 1.55\text{ ng g}^{-1}\text{ dw}$  (mean  $\pm$  standard deviation,  $n = 3$ ), respectively. The concentration of BDE-209 in the contaminated diet was considered to be ecologically relevant and representative of high levels (i.e.  $\sim 10\times$  higher the average concentrations) of this congener contamination found in marine species inhabiting contaminated coastal areas (e.g. Noyes and Stapleton, 2014; Airaksinen et al., 2015; Cruz et al., 2015).

## 2.3. Biochemical analyses

Dry matter, ash, total lipids, crude protein and gross energy were determined in the diet and fish body according to the Association of Official Analytical Chemists methods (AOAC, 2005). The contents of crude protein and gross energy were also determined in the faeces. Briefly, moisture was determined by drying the sample overnight at  $105^\circ\text{C}$  (lab heater, P-Selecta 207, Portugal); ash by incineration in a muffle furnace at  $500^\circ\text{C}$  for 16 h (Heraeus Hanau, TYP. MR170, Germany); and total lipids were determined with the Soxhlet extraction method using ethyl ether. Crude protein ( $N \times 6.25$ ) was analysed using the Dumas method, in a nitrogen analyser (LECO, Model FP-528, St Leco Corporation, St. Joseph, MI, USA) calibrated with EDTA. Gross energy was determined by combustion in an adiabatic bomb calorimeter (Werke C2000, IKA, Staufen, Germany) calibrated with benzoic acid. All chemical analyses were carried out in duplicate.

## 2.4. Excretion

### 2.4.1. Determination of faecal losses

During the 56 days of each experiment, faeces were collected daily by siphoning before feeding, oven-dried at  $65^\circ\text{C}$ , weighted, homogenized and stored at  $-20^\circ\text{C}$  for analysis of energy and nitrogen content. The faeces collection from each tank were pooled ( $n = 3$  per treatment) and measured by combustion in an adiabatic bomb calorimeter (Werke C2000, IKA, Staufen, Germany).

### 2.4.2. Determination of nitrogenous losses (non-faecal excretion)

Energy loss through excreted nitrogenous compounds (ammonia-N) was determined at the end of the experiment using 5 fish enclosed in  $500\text{ mL}$  glass bottles for 1 h for each treatment. Controls without fish (two blanks per treatment) were also used and ammonia levels were measured to evaluate the potential loss of nitrogenous compounds through bacterial action or diffusion in the experimental tanks. The bottles were filled with oxygen saturated seawater and sealed without any air bubbles. Trials were carried out in the same conditions (temperature, pH and salinity) of each treatment. The seawater was filtered and frozen in  $20\text{ mL}$  plastic flasks for ammonia quantification. Ammonia concentration was determined according to Berthelot method (Grasshoff, 1983). Samples were treated with alkaline citrate, sodium hypochloride and fenol in the presence of sodium nitroprussiate which catalyses the reaction. The blue colour formed by indofenol plus ammonia reaction was measured at  $630\text{ nm}$ . Ammonia excretion rate (AER; expressed as  $\text{mg kg}^{-1}\text{ dw L}^{-1}\text{ h}^{-1}$ ) was calculated as follows:

$$\text{AER} = \Delta[\text{NH}_4^+] \times \text{VH}_2\text{O}/\text{DW}/\Delta\text{T},$$

where  $\Delta[\text{NH}_4^+]$  is the difference between the ammonia levels in the sample and blank,  $\text{VH}_2\text{O}$  is the volume of water (L) in the glass bottle, DW is the fish dry weight, and  $\Delta\text{T}$  is the time duration of the trial (hours).

## 2.5. Metabolism

Oxygen consumption rates (routine metabolic rates; RMRs) were determined according to previously established methods (Rosa and Seibel, 2008, 2010; Anacleto et al., 2014) with some modifications. At

day 56 of the experiment, respiration trials were conducted in an intermittent flow-through respirometer system to measure oxygen consumption rates and to determine RMRs. Respiration chambers were filled with seawater from the experimental tank to ensure the correct temperature, pH and dissolved oxygen levels and placed in thermostable water baths (Lauda, Lauda-Königshofen, Germany) in order to control the temperature. The seawater parameters were controlled using a WTW hand-held Meter Multi 350i compact (WTW, Wissenschaftlich-Technische Werkstätten, Weilheim, Germany) in order to ensure that conditions were maintained under the correct parameters levels. Oxygen levels within the respirometers remained above 70% air saturation at all times. Juvenile fish were fasted overnight before being placed in the respirometers and allowed to acclimate (10 min) before data collection for stabilisation of oxygen uptake rates. Individual fish from each experimental tank were individually weighted (range: 5.3–13.4 g wet weight; ww) and placed into separate cylindrical respirometers (7.1 cm diameter, 15 cm length, 15 cm width, 14 cm depth, 545 mL volume; Loligo Systems, Tjele, Denmark), with a removable lid for introduction and removal of the fish. One of the chambers was used as control for each treatment and the oxygen consumption observed in this blank chamber was then subtracted from the values obtained in chambers containing fish. Six fish were used from each experimental tank/treatment. Respirometers were connected to an automated flush pump that flushed the respirometers with air-saturated seawater until attaining initial oxygen levels. The concentration of dissolved oxygen in the respirometers were monitored and recorded with an fibre-optic system and contactless oxygen sensor spots (FireSting O<sub>2</sub>, PyroScience, GmbH, Aachen, Germany) using the Pyro Oxygen Logger software. Continuous monitoring of oxygen concentration in the chambers was performed visually through real-time graphics on the computer monitor. The respirometers with fish were covered with a black bag to reduce light levels and disturbance. Calibrations were performed with seawater containing 100% air saturation representing full oxygen saturation. Oxygen consumption rates were determined according to the decline in oxygen concentration in the respirometers between flush cycles. Intermittent flow enables measurements to be taken over an extended period of time without depleting dissolved oxygen concentrations in the respirometer below normoxic concentrations. Fish were returned to their respective holding tanks following respirometry. Respirometers were cleaned regularly to ensure that background microbial respiration remained negligible. Metabolic data were analysed after importing the text file from the FireSting O<sub>2</sub> software into Excel 2013. Linear regressions between water oxygen concentration and time were made for each measurement period and the slopes derived from the regressions were used to calculate routine metabolic rate (RMR, mg O<sub>2</sub> kg<sup>-1</sup> ww h<sup>-1</sup>) according to the following equation:

$$RMR = (b \times V \times t) / W$$

where *b* is the slope of the oxygen reduction curve (minus the blank control), *V* is the volume of the respirometer minus the volume of the fish assuming that 1 kg of seawater is equivalent to 1 kg of fish (in L), *t* is the time interval over which O<sub>2</sub> is assessed (in hours) and *W* is the wet weight of fish (in kg). Additionally, O:N ratios were calculated by dividing the amount (atoms) of oxygen consumed by the nitrogen excreted, being used to estimate what substrates (carbohydrate, protein and lipid) the organisms use for metabolism (Babarro et al., 2000):

$$O : N = (RMR/16)/(AER/17)$$

## 2.6. Data calculation

### 2.6.1. Survival and animal fitness

Survival rate (SR, %) was calculated according to the following equation:

$$SR = \text{final number of fish} / \text{initial number of fish} \times 100$$

The Fulton's condition index (K) was directly calculated from the biometric data to determine fish condition, according to the formula (Ricker, 1975):

$$K = TW / TL^3 \times 100$$

where TW is the fish total weight (g wet mass) and TL is the total length (cm).

The relationship between fish total weight and the respective organ weight was calculated for the liver (i.e. hepatosomatic index, HSI) and the viscera (i.e. viscerosomatic index, VSI), in order to provide information on liver and viscera condition, using the following equations:

$$HSI (\%) = \text{liver wet weight} / \text{total fish wet weight} (g) \times 100$$

$$VSI (\%) = \text{viscera wet weight} (g) / \text{total fish wet weight} (g) \times 100$$

### 2.6.2. Growth performance

Relative growth rate (RGR, %), specific growth rate in terms of protein (SGRp, % day<sup>-1</sup>) and energy (SGRe, % day<sup>-1</sup>), feed intake (FI, % body weight day<sup>-1</sup>) and feed efficiency (FE) were calculated according to the following equations (Fang et al., 2010):

$$RGR = [(W2 - W1) / W1] \times 100$$

$$SGRp = [(\ln P2 - \ln P1) / t] \times 100$$

$$SGRe = [(\ln E2 - \ln E1) / t] \times 100$$

$$FI = [F / (t \times (W2 + W1/2))] \times 100$$

$$FE = (W2 - W1) / F$$

where W2 and W1 were the final and initial body weight of the fish, respectively, P2 and P1 were final and initial protein content, respectively, E2 and E1 were final and initial energy content, respectively, *t* was the duration of the experiment (days) and *F* was the dry weight of feed intake during the experimental period (g).

### 2.6.3. Energy budgets

Energy budgets were calculated according the model proposed by Carfoot (1987):

$$C = G + F + U + R,$$

where *C* is energy consumed, *G* is energy accumulated for growth, *F* is the energy loss in faeces, *U* is the energy loss as ammonia excretion; and *R* is the energy loss through respiration. Value of *G* in the budget equation was calculated as follows:

$$G = (W2 \times E2) - (W1 \times E1)$$

where W2 and W1 are the final and initial wet body weights of fish (g), respectively; and E2 and E1 are final and initial energy contents of the fish (J g<sup>-1</sup>), respectively.

The energy values of *F* were determined using an adiabatic bomb calorimeter (Werke C2000, IKA, Staufen, Germany). For determination of value of *U*, ammonia-N levels (AER) were converted into energy by multiplying the coefficient of 24.83 kJ g<sup>-1</sup> N (Elliott, 1976). The oxygen consumption was converted to energy (value of *R*) using a conversion factor of 13.84 J mg<sup>-1</sup> O<sub>2</sub> (442.88 J mmol<sup>-1</sup> O<sub>2</sub>) (Guinea and Fernandez, 1997).

The assimilation efficiency (K<sub>1</sub>, %) and net growth efficiency (K<sub>2</sub>, %) were calculated using the following equations (Ye et al., 2009):

$$K1 = [(G + R) / (G + R + U)] \times 100$$

$$K2 = [G / (G + R)] \times 100$$

## 2.7. Statistical analysis

All data were checked for normality of distribution and homogeneity of variances using Kolmogorov–Smirnov and Levene's tests,



**Table 1**

Survival, initial wet weight, final wet weight, total length, Fulton's condition index, hepatosomatic and visceral indices, feed intake and feed efficiency of juvenile white seabream (*Diplodus sargus*) in the four treatments after 56 days of experiment.

	19 °C_Control	19 °C_BDE-209 exposure	24 °C_Control	24 °C_BDE-209 exposure
Survival (%)	90.0 ± 0.0	88.2 ± 10.2	90.2 ± 9.0	94.1 ± 5.9
Wi (g)	5.07 ± 1.41 <sup>ab</sup>	4.50 ± 0.80 <sup>ab</sup>	3.93 ± 1.09 <sup>b</sup>	5.95 ± 1.25 <sup>a</sup>
Wf (g)	9.61 ± 2.29 <sup>ab</sup>	7.05 ± 1.54 <sup>b</sup>	9.88 ± 2.07 <sup>ab</sup>	10.26 ± 2.99 <sup>a</sup>
TL (cm)	8.04 ± 0.66 <sup>a</sup>	7.23 ± 0.44 <sup>b</sup>	8.03 ± 0.55 <sup>a</sup>	8.03 ± 0.74 <sup>a</sup>
K	1.82 ± 0.11	1.84 ± 0.14	1.88 ± 0.08	1.93 ± 0.14 <sup>†</sup>
HSI	1.30 ± 0.56	1.25 ± 0.33	0.90 ± 0.15 <sup>†</sup>	1.21 ± 0.29
VSI	6.87 ± 0.81 <sup>a,*</sup>	6.27 ± 0.91 <sup>ab</sup>	5.57 ± 1.10 <sup>b</sup>	6.05 ± 0.87 <sup>ab</sup>
FI (g g <sup>-1</sup> day <sup>-1</sup> )	0.47 ± 0.10	0.48 ± 0.04	0.52 ± 0.10	0.43 ± 0.09
FE	0.72 ± 0.23 <sup>b</sup>	0.70 ± 0.33 <sup>b</sup>	1.15 ± 0.14 <sup>b</sup>	1.47 ± 0.32 <sup>a</sup>

Values (mean ± SD; n = 9) with different letters (a-b) in the same row indicate significant differences between treatments (ANOVA test, p < 0.05). Asterisk (\*) represent significant differences between initial K, HSI and VSI and each treatment (t-student, p < 0.05). Abbreviations: 19 °C\_Control – seawater temperature set at 19 °C and animals fed with the control diet; 19 °C\_BDE-209 exposure – seawater temperature set at 19 °C and animals fed with BDE-209 contaminated diet; 24 °C\_Control – seawater temperature set at 24 °C and animals fed with the control diet; 24 °C\_BDE-209 exposure – seawater temperature set at 24 °C and animals fed with BDE-209 contaminated diet; Wi – Initial wet weight; Wf – Final wet weight; TL – Total length; K – Fulton's condition index; HSI – hepatosomatic index; VSI – viscerosomatic index; FI – Feed intake; FE – Feed efficiency.

respectively. Two-way analysis of variance (ANOVA) was used to analyse the effects of temperature and exposure to dietary BDE-209 on energy budget parameters. One-way ANOVA, followed by post-hoc tests (Tukey HSD) or non-parametric multiple comparison tests (Dunn's) were performed to analyse differences in indices (K, HSI, VSI, SR, RGR, SGR, FI and FE) and energy budget parameters (C, G, F, U and R) among treatments. The correlation between variables (energy budget parameters and other indexes, temperature and BDE-209 contaminated diet) was tested by Pearson's correlation coefficients. Differences were considered significant at a probability levels of p < 0.05, using STATISTICA™ software (Version 10.0, StatSoft Inc., Tulsa, Oklahoma, USA). All data were expressed as mean ± standard deviation (mean ± SD).

### 3. Results

#### 3.1. Survival and fitness

Throughout the 56-day experiment, the survival rate of juvenile white seabream ranged between 88.2% and 94.1% in treatments 19 °C\_BDE-209 exposure and 24 °C\_BDE-209 exposure, respectively (Table 1). There was no significant differences in survival rate between the four treatments (p > 0.05). The mortality rate observed in all treatments was mainly caused by cannibalism as this species demonstrated a normal aggressive behaviour.

Despite no significant differences were observed in Fulton's

**Table 2**

Effects of warming and BDE-209 contamination on biochemical composition of juvenile white seabream (*Diplodus sargus*) after 56 days of the experiment.

	Initial	19 °C_Control	19 °C_BDE-209 exposure	24 °C_Control	24 °C_BDE-209 exposure
Moisture (%)	72.67 ± 0.23	70.81 ± 1.45	72.05 ± 1.39	71.64 ± 0.26 <sup>†</sup>	71.46 ± 0.59 <sup>†</sup>
Protein (%)	54.78 ± 1.63	50.85 ± 0.39 <sup>b,*</sup>	50.98 ± 0.65 <sup>b,*</sup>	54.63 ± 0.93 <sup>a</sup>	49.17 ± 0.16 <sup>b,*</sup>
Fat (%)	19.90 ± 0.06	26.10 ± 0.62 <sup>a,*</sup>	24.66 ± 0.13 <sup>ab,*</sup>	23.45 ± 1.10 <sup>bc,*</sup>	21.63 ± 0.38 <sup>c,*</sup>
Ash (%)	18.55 ± 0.39	16.66 ± 0.34 <sup>b,*</sup>	17.03 ± 0.15 <sup>ab,*</sup>	17.84 ± 0.40 <sup>ab,*</sup>	18.30 ± 0.48 <sup>a</sup>
Gross energy (kJ g <sup>-1</sup> )	18.87 ± 0.39	20.48 ± 0.33 <sup>†</sup>	20.21 ± 0.24 <sup>†</sup>	19.70 ± 0.18 <sup>†</sup>	20.29 ± 0.32 <sup>†</sup>

Values (mean ± SD; n = 3) correspond to the percentage of dry body weight. Different letters in the same row indicate significant differences between the four treatments (ANOVA test, p < 0.05). Asterisk (\*) represent significant differences between initial biochemical composition and each treatment (t-student, p < 0.05). Abbreviations: 19 °C\_Control – seawater temperature set at 19 °C and animals fed with the control diet; 19 °C\_BDE-209 exposure – seawater temperature set at 19 °C and animals fed with BDE-209 contaminated diet; 24 °C\_Control – seawater temperature set at 24 °C and animals fed with the control diet; 24 °C\_BDE-209 exposure – seawater temperature set at 24 °C and animals fed with BDE-209 contaminated diet.

condition (K) and hepatosomatic index (HSI) between the four treatments, the viscerosomatic index (VSI) was significantly changed by temperature (two-way ANOVA, F = 6.04, p < 0.05; Table A.2), i.e. fish exposed to control conditions (i.e. 19 °C\_Control) showed significantly higher VSI (6.87 ± 0.81) than fish subjected to warmer temperatures (i.e. 24 °C\_Control; 5.57 ± 1.10; F = 3.06, p < 0.05). Comparing to the initial values at the beginning of the experiment, significant differences were also observed for K at 24 °C\_BDE-209 exposure (p < 0.05), HSI at 24 °C\_Control and VSI at 19 °C\_Control (p < 0.01; Table 1).


#### 3.2. Biochemical composition

Moisture, protein, fat, ash and gross energy contents in the body of juvenile white seabream exposed to the different treatments are listed in Table 2. Fish body protein content was affected by both temperature and BDE-209 contaminated diet (two-way ANOVA, F = 6.23, p < 0.05 and F = 45.95, p < 0.001, respectively; Table A.3) as well as fish fat content (F = 44.86, p < 0.001 and F = 14.65, p < 0.01, respectively). Fish ash was only affected by temperature (F = 22.83, p < 0.01). No significant differences were observed between moisture and gross energy of the fish between the different treatments (p > 0.05). Yet, there was a significant interaction between temperature and BDE-209 contaminated diet in gross energy and also in protein contents (F = 6.03, p < 0.05 and F = 50.50, p < 0.001, respectively). In comparison with initial values of biochemical composition, significant differences were detected after 56 days of experiment (p < 0.05), except for moisture in both treatments at 19 °C, for protein in treatment 24 °C\_Control and for ash in treatment 24 °C\_BDE-209 exposure. Moisture, protein and ash contents significantly decreased after 56 days of experiment in each treatment, whereas fat and gross energy significantly increased (p < 0.05).

#### 3.3. Growth


Table 1 shows that significant differences were registered in the initial and final body weights of animals among treatments. Fish subjected to warming conditions and BDE-209 exposure (i.e. 24 °C\_BDE-209 exposure) obtained the highest body weights (Wi = 5.95 ± 1.25 g, F = 4.67, p < 0.01; Wf = 10.26 ± 2.99 g, F = 3.67, p < 0.05), whereas fish exposed to control temperature and BDE-209 exposure (i.e. 19 °C\_BDE-209 exposure) showed the lowest total length (TL = 7.23 ± 0.44 cm, F = 3.93, p < 0.05). A strong positive correlation was found between fish total weight and length (r = 0.95; p < 0.001; data not shown). The two-way ANOVA analysis also showed that temperature significantly affected final fish weights (F = 5.23, p < 0.05) and a significant interaction was registered between temperature and BDE-209 contaminated diet in the initial body weight (F = 9.83, p < 0.01; Table A.2).

Feed intake (FI) and specific growth rate in terms of energy and protein (SGRe and SGRp, respectively) did not show significant



**Fig. 1.** A) Relative growth rate (RGR, %) and B, C) specific growth rate in terms of energy and protein (SGRe and SGRp, respectively, % day<sup>-1</sup>) in juvenile white seabream (*Diplodus sargus*) after 56 days of exposure to warming and contaminant (mean ± SD; n = 9). Different letters indicate significant differences between treatments (ANOVA test, p < 0.05). Abbreviations: 19°C\_Control – seawater temperature set at 19 °C and animals fed with the control diet; 19°C\_BDE-209 exposure – seawater temperature set at 19 °C and animals fed with BDE-209 contaminated diet; 24°C\_Control – seawater temperature set at 24 °C and animals fed with the control diet; 24°C\_BDE-209 exposure – seawater temperature set at 24 °C and animals fed with BDE-209 contaminated diet.

differences within treatments (Table 1 and Fig. 1B and C). However, FE was significantly affected by temperature and BDE-209 contaminated diet (two-way ANOVA, F = 71.16, p < 0.001 and F = 23.18, p < 0.001, respectively; Table A.2) and RGR was only significantly affected by temperature (F = 27.71, p < 0.01; Table A.2). Moreover, a significant interaction between temperature and BDE-209 contaminated diet occurred only for FE (F = 24.52, p < 0.001; Table A.2). Fish exposed to warmer temperatures and contaminated diet (i.e. 24°C\_BDE-209 exposure) revealed significantly higher FE (1.47 ± 0.32; Table 1) and RGR levels (107.1 ± 18.0; Fig. 1A).



**Fig. 2.** Effects of warming and BDE-209 contamination on A) ammonia excretion rate (AER; mg NH<sub>4</sub> g<sup>-1</sup> dw L<sup>-1</sup> h<sup>-1</sup>), B) routine metabolic rate (RMR; mg O<sub>2</sub> kg<sup>-1</sup> ww h<sup>-1</sup>) and C) O:N ratio of juvenile white seabream (*Diplodus sargus*) after 56 days of exposure (mean ± SD; n = 6). Different letters indicate significant differences between treatments (ANOVA test, p < 0.01). Abbreviations: 19°C\_Control – seawater temperature set at 19 °C and animals fed with the control diet; 19°C\_BDE-209 exposure – seawater temperature set at 19 °C and animals fed with BDE-209 contaminated diet; 24°C\_Control – seawater temperature set at 24 °C and animals fed with the control diet; 24°C\_BDE-209 exposure – seawater temperature set at 24 °C and animals fed with BDE-209 contaminated diet.

### 3.4. Excretion and metabolism

Ammonia excretion rate (AER) was significantly affected by temperature and BDE-209 contaminated diet (two-way ANOVA, F = 5.34, p < 0.05 and F = 22.30, p < 0.001, respectively; Table A.2). Indeed, fish exposed to warming conditions (i.e. 24°C\_Control) showed significantly higher AER (0.076 ± 0.021 mg NH<sub>4</sub> g<sup>-1</sup> dw L<sup>-1</sup> h<sup>-1</sup>), whereas fish subjected to contaminated diet (i.e. 19°C\_BDE-209 exposure and 24°C\_BDE-209 exposure) presented significantly lower AER (0.084 ± 0.007 and 0.111 ± 0.023 mg NH<sub>4</sub> g<sup>-1</sup> dw L<sup>-1</sup> h<sup>-1</sup>,

respectively;  $F = 9.22$ ,  $p < 0.01$ ; Fig. 2A).

Regarding routine metabolic rates (RMRs), fish subjected to control temperature and BDE-209 exposure displayed significantly lower levels ( $464 \pm 76 \text{ mg O}_2 \text{ kg}^{-1} \text{ ww h}^{-1}$ ; Fig. 2B) than the other treatments ( $p < 0.01$ ). The two-way ANOVA showed that RMRs of juvenile white seabream were significantly affected by both temperature and BDE-209 contaminated diet ( $F = 15.96$ ,  $p < 0.001$  and  $F = 17.60$ ,  $p < 0.001$ ; Table A.2) and there was also a significant interaction between temperature and BDE-209 contaminated diet ( $F = 6.48$ ,  $p < 0.05$ ).

O:N ratios of the juvenile white seabream subjected to different treatments are shown in Fig. 2C. The O:N ratios were significantly lower in control diet treatments ( $15.23 \pm 1.52$  and  $15.83 \pm 1.04$  in  $19^\circ\text{C\_Control}$  and  $24^\circ\text{C\_Control}$ , respectively) than in contaminated diet treatments ( $19.35 \pm 1.10$  and  $21.22 \pm 0.90$  in  $19^\circ\text{C\_BDE-209}$  exposure and  $24^\circ\text{C\_BDE-209}$  exposure, respectively;  $F = 22.07$ ,  $p < 0.001$ ). The two-way ANOVA revealed that O:N ratio of juvenile white seabream was only significantly affected by BDE-209 exposure ( $F = 60.34$ ,  $p < 0.001$ ; Table A.2).

### 3.5. Energy budget

Energy intake from feed and energy allocation in percentage of total energy intake is presented in Fig. 3A and B. Energy intake showed significant differences within treatments ( $p < 0.001$ ): warming conditions and BDE-209 contaminated diet exposure (i.e.  $24^\circ\text{C\_BDE-209}$  exposure) promoted higher energy acquired from feed ( $992 \pm 32 \text{ J g}^{-1} \text{ day}^{-1}$ ), whereas treatments with control temperature (i.e.  $19^\circ\text{C\_BDE-209}$  exposure and  $19^\circ\text{C\_Control}$ ) showed the lowest levels of energy intake ( $562 \pm 5$  and  $854 \pm 2 \text{ J g}^{-1} \text{ day}^{-1}$ , respectively).

Regarding energy distribution, the highest fraction of energy was allocated to growth (G) in all treatments (between 56.0% and 67.8%; Fig. 3B), followed by respiration (R; around 25–26%) and excretion through faeces (F; between 4.3% and 16.0%) and ammonia (U; between 2.3% and 3.3%). Growth energy was significantly higher under warming and BDE-209 exposure via diet (i.e.  $24^\circ\text{C\_BDE-209}$  exposure =  $67.8 \pm 1.4\%$ ; Fig. 3B), compared to the control temperature and BDE-209 exposure via diet (i.e.  $19^\circ\text{C\_BDE-209}$  exposure;  $56.0 \pm 1.4\%$ ;  $F = 29.85$ ,  $p < 0.01$ ). On the other hand, energy loss via faeces (F) exhibited an opposite trend to growth, with the latter treatment showing the highest percentage ( $19^\circ\text{C\_BDE-209}$  exposure =  $16.0 \pm 0.9\%$ ; Fig. 3B), while significant lower percentage was observed under  $24^\circ\text{C\_BDE-209}$  exposure ( $4.3 \pm 0.1\%$ ;  $F = 201.47$ ,  $p < 0.0001$ ). Non-significant differences between treatments were observed for energy lost through ammonia excretion and respiration ( $p > 0.05$ ).

In general, energy intake and energy distribution were significantly influenced by temperature and emerging BDE-209 exposure via diet ( $p < 0.05$ ; Table A.3), except respiration that was not influenced by temperature ( $p > 0.05$ ). Concerning assimilation and net growth efficiencies ( $K_1$  and  $K_2$ , respectively), no significant differences in  $K_2$  were detected between any treatment ( $p > 0.05$ ), whereas  $K_1$  was only significantly lower under  $19^\circ\text{C\_BDE-209}$  exposure ( $88.9\%$ ;  $F = 109.1$ ,  $p < 0.001$ ; Fig. 3C).

## 4. Discussion

Several studies have assessed the effects of temperature on the energy budget of different fish species, whereas others only focused on the effects of chemical contaminants under a climate change context, without ever reporting the effects on energy budgets. Moreover, the interactive effects of these environmental stressors on fish bioenergetics are poorly known and require further understanding. Therefore, this study aimed to attenuate this knowledge gap by providing the first empirical evidence of the combined impacts of warming and contamination on fish energy budget.

Since temperature is recognized as the most important abiotic factor




Fig. 3. Effects of warming and BDE-209 contamination on A) energy intake from feed ( $C$ ;  $\text{J g}^{-1} \text{ day}^{-1}$ ), B) energy allocation (%), C) assimilation efficiency ( $K_1$ ) and net growth efficiency ( $K_2$ ) of juvenile white seabream (*Diplodus sargus*) after 56 days of the experiment. Values (mean  $\pm$  SD;  $n = 6$ ) are represented as the percentage of dry body weight. Different letters indicate significant differences between the treatments (ANOVA test,  $p < 0.05$ ). Abbreviations: C - energy intake; G - energy deposit for growth; F - energy loss through faeces; U - energy loss through ammonia excretion; R, energy loss through respiration;  $K_1$  - assimilation efficiency;  $K_2$  - net growth efficiency;  $19^\circ\text{C\_Control}$  - seawater temperature set at  $19^\circ\text{C}$  and animals fed with the control diet;  $19^\circ\text{C\_BDE-209}$  exposure - seawater temperature set at  $19^\circ\text{C}$  and animals fed with BDE-209 contaminated diet;  $24^\circ\text{C\_Control}$  - seawater temperature set at  $24^\circ\text{C}$  and animals fed with the control diet;  $24^\circ\text{C\_BDE-209}$  exposure - seawater temperature set at  $24^\circ\text{C}$  and animals fed with BDE-209 contaminated diet.

affecting physiological functioning of most marine (poikilothermic) organisms (i.e. body temperature varies over time with environmental temperature), its role on fish energy budget has also been intensively investigated, as in *Sparus aurata* (Requena et al., 1997), *Trematomus bernacchii* (Sandersfeld et al., 2015), *Cynoglossus semilaevis* (Fang et al., 2010), *Oncorhynchus tshawytscha*, *Salvelinus namaycush*, *Oncorhynchus mykiss* (Kao et al., 2015), *Oreochromis niloticus* (Xie et al., 2011), *Rachycentron canadum* (Sun et al., 2006; Sun and Chen, 2009) and *Gadus morhua* (Kreiss et al., 2015). However, the conclusions on the influence of temperature on energy budget of species were not consistent among these studies.

When an environmental change occurs (ex. warming), an organism needs to perform metabolic adjustments, which will be stressful and have direct consequences on its fitness that is influenced by the energy budget. Here, despite survival, fish condition (K) and hepatosomatic index (HSI) were not significantly affected by temperature, viscerosomatic index (VSI) decreased significantly with warming (Table 1), which is correlated with fat content. Indeed, other authors demonstrate that VSI and fat content increase at low temperature, which may be linked to a deposition of liver glycogen and lipids (Couto et al., 2008; Sandersfeld et al., 2015; Huang et al., 2016). This finding it is not surprising taking into account that temperature can influence fish body composition, by affecting the stability of structural components, particularly lipids and proteins (Van Ham et al., 2003; Fang et al., 2010). Similar results were reported by Brodte et al. (2006) for lipid content in the Antarctic eelpout. Such response to warming may be led by a shift from lipid- to carbohydrate-based metabolism (Windisch et al., 2011). We can also argue that the lower content of fat in fish under warming (Table 2) maybe also a consequence of enhanced metabolic processes associated to growth (García-Guerrero et al., 2003). In fact, protein and ash contents increased with warming in juvenile white seabream. It is worth noting that the use of the protein fraction as an energy source for the maintenance of animal metabolic needs only takes place when carbohydrate and lipid reserves have already been greatly depleted (Barber and Blake, 1981). Proteins have an important physiological role in the supply of structural elements and in the catalysis of metabolic reactions (Dittrich, 1991). Regarding gross energy, warming and BDE-209 exposure did not affect the content of *D. sargus*.

Despite no differences were observed in specific growth rate in terms of protein and energy (SGRp and SRGe, respectively; Fig. 1B and C), temperature had a significant positive effect on fish growth, namely in terms of an increase in wet weight and relative growth rate (Table 1 and Fig. 1A). It is well known that temperature can affect fish growth directly by controlling feed consumption, nutrient requirements and food passage time (Smith, 1989). While increasing temperature may elicit negative effects on growth due to higher energy cost for maintenance metabolism, may also elicit positive ones due to higher food consumption and digestion rates (Xie and Sun, 1992, 1993). Sandersfeld et al. (2015) observed a reduction in mass growth (84%) in Antarctic fish *Trematomus bernacchii* at 2 and 4 °C when compared with the control group at 0 °C, which can be explained by reduced food assimilation rates at warmer temperatures. Within a suitable temperature range, growth rate increases with increasing temperature, but when the temperature reaches the upper extreme of the tolerated range, usually the opposite happens (Hasan and Macintosh, 1991). Regarding *D. sargus*, its wide continuous distribution range in temperate/subtropical areas indicates that this species tolerates water temperatures until 34 °C, but the preferred temperatures range between 15 and 17 °C, which corresponds to the spawning period that generally occurs between March to June (Morato et al., 2003).

Concerning feed efficiency (FE), there was a significant increase under warming (Table 1), which is in accordance with Guerreiro et al. (2012) that also obtain higher FE at 22 °C than at 16 °C, suggesting that this temperature can be nearer the optimum temperature for the species as FE is usually maximized at temperatures near the optimum for maximum growth of the species (Ármason et al., 2009). Although it is well established that fish eat to meet their energetic demands (Bureau et al., 2002) within limits, they may also adjust feed intake to meet protein requirement (Peres and Oliva-Teles, 1999). On the other hand, our results are contradictory with other authors that demonstrated a decrease in FE when temperature increased (e.g. *Rachycentron canadum*; Sun et al., 2006). Other study performed on young grass carp (Cui et al., 1995) showed that temperature did not significantly affect FE. This discrepancy in the results probably resulted from interspecific differences, or from the reason that the designed temperature in the experiment of a given species was not beyond its suitable temperature range.

In relation to the excretion and metabolic rates (AER and RMR, respectively), our results also showed that both increased with warming conditions in juvenile *D. sargus*. The higher levels of AER observed with increased temperature ( $0.176 \pm 0.021 \text{ mg NH}_4 \text{ g}^{-1} \text{ dwL}^{-1} \text{ h}^{-1}$ ; Fig. 2A) are consistent with previous findings for other fish species (Tidwell et al., 2003; Leung et al., 1999). Also, Forsberg and Summerfelt (1992) have shown that the AER in *Stizostedion vitreum vitreum* juveniles varies between groups exposed to 20 and 25 °C ( $30.1$  and  $45.2 \text{ mg NH}_4 \text{ kg}^{-1} \text{ h}^{-1}$ , respectively). In comparison to *D. sargus*, AER was slightly higher, varying between  $41.7$  and  $50.0 \text{ mg NH}_4 \text{ g}^{-1} \text{ h}^{-1}$  at 19 and 24 °C, respectively (Fig. 2A). This can be due to the fact that at higher temperatures, increases in metabolic rates are partially attained via the transamination and deamination of the amino acids ingested in the diet, resulting in the excretion of ammonium and the release of left-over carbon used as an energy source (Forsberg and Summerfelt, 1992).

The increase in RMR of juvenile *D. sargus* with warming was observed in both control and contaminated treatments, although was only significant under exposure to the flame retardant (i.e.  $463.8 \pm 75.8 \text{ mg O}_2 \text{ kg}^{-1} \text{ ww h}^{-1}$  at 19 °C to  $634.6 \pm 99.5 \text{ mg O}_2 \text{ kg}^{-1} \text{ ww h}^{-1}$  at 24 °C; Fig. 2B and Table A.2). Since temperature directly affects the rate of all biological processes, this positive trend between the two variables is expected and consistent with the results of previous studies for other fish species, such as *Sparus aurata* (Requena et al., 1997). Indeed, each species has a thermal optimum at which it functions are the best, and increasing temperatures beyond that point will cause metabolic rates to decline (Pang et al., 2015). For example, Chen et al. (2006) found that between 6 and 34 °C, RMR of juvenile common carp drastically decreased above 30 °C. Moreover, species also has an upper and lower lethal temperatures or critical thermal maxima or minima (CT<sub>max</sub> and CT<sub>min</sub>, respectively), beyond which metabolic functions cannot continue. Therefore, the concept of oxygen and capacity-limited thermal tolerance (OCLTT) demonstrated by Pörtner and Knust (2007) aims to explain this effect of temperature on body functioning. With water temperatures gradually rising as climate change progresses, temperatures may move farther away from species' thermal optima and closer to their thermal maxima, creating unfavourable environmental conditions for many species (Iannucci, 2015). In this study, it is apparent that the temperature have not reached maximum enough (at least 24 °C) to make their metabolic damage to juvenile *D. sargus*.

The O:N ratio is an effective index for assessing the contribution of protein catabolism to total metabolism and can provide indices of balance between the catabolism rates of proteins, carbohydrates, and lipid substrates in animal tissues (Jadhav et al., 2012). This parameter is affected by both endogenous physiological status and exogenous environmental factors (Rocha et al., 2005). High O:N ratio (> 30) points out catabolism of carbohydrates and lipids (Bayne, 1976), while low O:N ratio (< 30) rather points out protein catabolism, i.e. conditions of alimentary deficiency (Mace and Ansell, 1982). In this study, results showed a prevalence of protein-dominated catabolism in all treatments (between  $15.22 \pm 1.52$  and  $21.22 \pm 0.90$ ; Fig. 2C). Similar values of O:N ratio were found by Lemos et al. (2006) and Rocha et al. (2005). On the other hand, Zheng et al. (2008) observed a change of O:N ratio from 13.12 to 20.91 at temperatures of 6, 10, 15 and 20 °C to 51.37 at 25 °C for juvenile *Müchthys müyü*, which is indicative of the change in the metabolic substrate (lipids and carbohydrates instead of proteins) that can be related to lower nitrogen excretion. Within this point of view, it is more cost-effective to catabolize fat rather than protein (1 g will release approximately 9.5 and 4.5 kcal of energy, respectively; Cook et al., 2000), being an adaptive strategy of fish with greater thermal tolerance. Interestingly, O:N ratio in the present study was only affected by BDE-209 exposure and not by warming (Table A.2), whereby fish subjected to BDE-209 contaminated diet exposure presented higher O:N ratio ( $19.35 \pm 1.10$  and  $21.22 \pm 0.90$  at 19 and 24 °C, respectively; Fig. 2C).



In this study, we obtained the following energy budget equations for juvenile white seabream for respective treatments: 100C = 62.6G + 9.4F + 2.5U + 25.5R (19 °C\_Control); 100C = 56.0G + 16.0F + 2.6U + 25.4R (19 °C\_BDE-209 exposure); 100C = 62.7G + 7.4F + 3.3U + 26.5R (24 °C\_Control); and 100C = 67.8G + 4.3F + 2.3U + 25.6R (24 °C\_BDE-209 exposure; Fig. 3B). In general, the higher proportion of food energy was allocated to growth (56.0–67.8%) and metabolism (25.4–26.5%), which is in accordance with several studies (e.g. Tang et al., 2003; Sandersfeld et al., 2015), but in inverse order of magnitude, i.e. higher energy channelled for metabolism, followed by growth. Generally, in most studies performed with fish, metabolism costs usually make up 50% of total energy expenditure and growth around 30% (Fang et al., 2010; Xie et al., 2011). Nevertheless, the animals used in this experiment were juvenile fish, which means that species do not reproduce before having reached a certain size. It is known that this species reaches sexual maturity at 2 years, with an approximate size of 17 cm, whereby the energy expenditure of studied animals is only channelled to growth and not yet to reproduction. Therefore, factors influencing energy allocation and thereby growth in this species are likely to have far-reaching consequences for life history. The obtained results also contradict research conducted by Tang et al. (2003) regarding to the excretion parameters, i.e. higher energy allocated to faeces (4.3–16.0%), followed by ammonia (2.3–3.3%; Fig. 3B). According to Sun et al. (2006), faeces and nitrogenous excretion only account for a small portion of food energy in carnivorous fish species and do not greatly influence the proportion of food energy allocated to growth, so it is the metabolism that plays an important role in influencing the proportion of energy intake allocated to growth. Other authors also detected opposite findings in relation to the percentages of energy allocation for juvenile fish species, namely *Cynoglossus semilaevis* (Fang et al., 2010), *Rachycentron canadum* (Sun et al., 2006) and *Trematomus bernacchii* (Sandersfeld et al., 2015). Besides temperature, other factors can influence the energy budget of an organism, such as feed type or composition, ease of finding and consuming food, experimental method, fish body size, current fish life stage, and environment-specific stressors such as predator density (Jobling, 1994; Xie et al., 1997; Tang et al., 2003; Yang et al., 2003). For example, Fang et al. (2010) indicated that the highest proportion for growth could be obtained at a ration below satiation or at satiation.

Concerning the flame retardant studied, BDE-209 is the most commonly used commercial PBDE, and is of particular concern due to its bioaccumulation and biomagnification capacity via aquatic food web (Noyes et al., 2011). Moreover, this abundant congener can be metabolised into lower brominated congeners with higher toxicity to aquatic organisms (Kuo et al., 2010). Studies have associated BDE-209 exposure to liver toxicity, reproductive toxicity, endocrine toxicity and potentially carcinogenic effects (Du et al., 2008). However, a limited number of published studies investigated the effects of BDE-209 dietary exposure only on growth of marine species. Zhang et al. (2013) observed acute toxic effects on the growth and inter-specific competition of marine bloom-forming microalgae. Sha et al. (2015) also detected a significant influence of this congener on the population growth and reproduction parameters of rotifer, *Brachionus plicatilis*. Other authors showed that BDE-209 through dietary exposure during 30 days even at low concentrations ( $2 \mu\text{g g}^{-1}$ ) affected negatively juvenile fish growth rates (*Coregonus clupeaformis*) by measuring the sagittal otolith increment width (Kuo et al., 2010). To our knowledge, this study was the first that measured all the energy budget parameters to evaluate the potential effects of this contaminant uptake. The presented results revealed that BDE-209 exposure during 56 days significantly affects the energy budget parameters (C, G, F, U and R) as well as the assimilation efficiency ( $K_1$ ; Table A.3). The combined conditions (i.e. 24 °C\_BDE-209 exposure) promoted higher energy acquired from food and also higher energy was particularly channelled to growth, whereas lower energy was conducted to excretion via faeces and ammonia losses (Fig. 3A and B).

## 5. Conclusions

In conclusion, the present study indicated that warming and BDE-209 contaminated diet exposure had significant effects on the energy budget of juvenile white seabream. The energy partitioned for metabolism varied around 26% under all treatments, and warmer temperatures generally promoted higher wet weight, RGR, protein and ash contents as well as lower VSI, while under flame retardant exposure promoted higher O:N ratio. Nevertheless, the combined effects of both stressors led to higher FE, AER and RMR and lower fat content. Further research combining other climate change stressors (e.g. acidification and hypoxia) and emerging chemical contaminants are needed to better understand and forecast the biological responses in the ocean of tomorrow.

## Acknowledgments

This study was supported by CERES project (European H2020 program; ref. 678193). Sparos Lda company for providing the fish feed, and the team from EPPA aquaculture pilot station (Olhão) for providing the juvenile fish for trials. The authors are also very grateful to Marta Nogueira, Teresa Gama Pereira and Margarida Saavedra from IPMA for their assistance with some technical matters. This study was supported by the Portuguese Foundation for Science and Technology (FCT) through a post-doc Grant of P.A. (SFRH/BPD/100728/2014), as well as the Ph.D. Grant of A.L.M. (SFRH/BD/103569/2014) and the contracts of A.M. and R.R. in the framework of the IF2013 and IF2014 programs, respectively.

## Competing interests

The authors declare that they have no actual or potential competing interests.

## Ethical statement

Fish trials were approved by the Ethical Committee of the Faculty of Sciences of the Lisbon University (ORBEA) and were conducted in accordance with legal regulations (EU Directive 2010/63).

## Funding source

The research leading to these results has received funding from European H2020 program under the CERES project (Ref. 678193). The Portuguese Foundation for Science and Technology (FCT) supported the post-doc Grant of P.A. (SFRH/BPD/100728/2014), as well as the Ph.D. Grant of A.L.M. (SFRH/BD/103569/2014), and the contracts of A.M. and R.R. in the framework of the IF2013 and IF2014 programs, respectively.

## Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.envres.2018.02.023>.

## References

- Airaksinen, R., Hallikainen, A., Rantakokko, P., Ruokojärvi, P., Vuorinen, P.J., Mannio, J., Kiviranta, H., 2015. Levels and congener profiles of PBDEs in edible Baltic, freshwater, and farmed fish in Finland. *Environ. Sci. Technol.* 49, 3851–3859.
- Alaee, M., Arias, P., Sjödin, A., Bergman, Å., 2003. An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release. *Environ. Int.* 29, 683–689.
- Anacleto, P., Maulvault, A.L., Lopes, V.M., Repolho, T., Diniz, M., Nunes, M.L., Marques, A., Rosa, R., 2014. Ecophysiology of native and alien-invasive clams in an ocean warming context. *Comp. Biochem. Physiol. Part A: Mol. Integr. Physiol.* 175, 28–37.
- AOAC, 2005. Official Methods of Analysis of AOAC International. In: Latimer, G., Horwitz, W. (Eds.), AOAC, 18th ed. AOAC International, Gaithersburg, pp. 3000.

- Árnason, J., Imsland, A.K., Gustavsson, A., Gunnarsson, S., Arnarson, I., Reynisson, H., Jonsson, A.F., Smaradottir, H., Thorarensen, H., 2009. Optimum feed formulation for Atlantic halibut (*Hippoglossus hippoglossus* L.): minimum protein content in diet for maximum growth. *Aquaculture* 291, 188–191.
- Babarro, J.M.F., Fernández-Reiriz, M.J., Labarta, U., 2000. Metabolism of the mussel *Mytilus galloprovincialis* from two origins in the Ria de Arousa (north-west Spain). *J. Mar. Biol. Assoc. U. K.* 80, 865–872.
- Barber, B.J., Blake, N.J., 1981. Energy storage and utilisation in relation to gametogenesis in *Argopecten irradians concentricus* (Say). *J. Exp. Mar. Biol. Ecol.* 52, 121–134.
- Bayne, B.L., 1976. In *Marine Mussels: Their Ecology and Physiology*. International Biological programme to Cambridge University Press, Cambridge, London, New York, Melbourne, pp. 159.
- Brodte, E., Knust, R., Pörtner, H.O., 2006. Temperature-dependent energy allocation to growth in Antarctic and boreal eelpout (Zoaridae). *Polar Biol.* 30, 95–107.
- Bureau, D.P., Kaushik, S.J., Cho, C.Y., 2002. Bioenergetics. In: Halver, J.E., Hardy, R.W. (Eds.), *Fish Nutrition*, 3rd ed. Academic Press, California, USA, pp. 1–59.
- Burel, C., Ruyet, P.L., Gaumet, F., Roux, A.L., Severe, A., Boeuf, G., 1996. Effects of temperature on growth and metabolism in juvenile turbot. *J. Fish. Biol.* 49, 678–692.
- Carfoot, T.H., 1987. *Animal Energetics*. Academic Press, New York, pp. 407–515.
- Chen, S.B., Fan, Z.T., Chen, W.X., 2006. The relationship of respiratory rate and oxygen consumption rate in common carp (*Cyprinus carpio haematopterus*) under different temperatures. *Chin. J. Northeast Agric. Univ.* 37(3), 352–356 (in Chinese with English abstract).
- Clarke, A., Johnston, N.M., 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. *J. Anim. Ecol.* 68, 893–905.
- Cook, J.T., Sutterlin, A.M., McNiven, M.A., 2000. Effect of food deprivation on oxygen consumption and body composition of growth-enhanced transgenic Atlantic salmon (*Salmo salar*). *Aquaculture* 188, 47–63.
- Costa, L.G., Giordano, G., 2011. Is decabromodiphenyl ether (BDE-209) a developmental neurotoxicant? *Neurotoxicology* 32, 9–24.
- Couto, A., Enes, P., Peres, H., Oliva-Teles, A., 2008. Effect of water temperature and dietary starch on growth and metabolic utilization of diets in gilthead sea bream (*Sparus aurata*) juveniles. *Comp. Biochem. Physiol. Part A: Mol. Integr. Physiol.* 151, 45–50.
- Cruz, R., Cunha, S.C., Casal, S., 2015. Brominated flame retardants and seafood safety: a review. *Environ. Int.* 77, 116–131.
- Cruz, R., Cunha, S.C., Marques, A., Casal, S., 2017. Polybrominated diphenyl ethers and metabolites – an analytical review on seafood occurrence. *Trends Anal. Chem.* 87, 129–144.
- Cui, Y.B., Chen, S.L., Wang, S.M., 1995. Effect of temperature on the energy budget of the grass carp, *Ctenopharyngodon idella* Val. *Oceanol. Limnol. Sin.* 26 (2), 169–174.
- Dittrich, B., 1991. Biochemical composition of the parasitic amphipod *Hyperia galba* in relation to age and starvation. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* 161, 441–449.
- Doney, S.C., Ruckelshaus, M., Duffy, J.E., Barry, J.P., Chan, F., English, C.A., Galindo, H.M., Grebmeier, J.M., Hollowed, A.B., Knowlton, N., Polovina, J., Rabalais, N.N., Sydeman, W.J., Talley, L.D., 2012. Climate change impacts on marine ecosystems. *Ann. Rev. Mar. Sci.* 4, 11–37.
- Du, H.Y., Zhu, L., Chen, Z.Z., Li, Y., Duan, Z.H., 2008. The progress of study on the toxicological effects of deca-brominated diphenyl ether. *J. Toxicol.* 22 (1), 50–52.
- Elliott, J.E., Wilson, L.K., Wakeford, B., 2005. Polybrominated diphenyl ether trends in eggs of marine and freshwater birds from British Columbia, Canada, 1979–2002. *Environ. Sci. Technol.* 39 (15), 5584–5591.
- Elliott, J.M., 1976. Energy losses in the waste products of brown trout (*Salmo trutta* L.). *J. Anim. Ecol.* 45, 561–580.
- Fang, J., Tian, X., Dong, S., 2010. The influence of water temperature and ration on the growth, body composition and energy budget of tongue sole (*Cynoglossus semilaevis*). *Aquaculture* 299, 106–114.
- Forsberg, J.A., Summerfelt, R.C., 1992. Effect of temperature on diel ammonia excretion of fingerling walleye. *Aquaculture* 102, 115–126.
- García-Guerrero, M., Villarreal, H., Racotta, I.S., 2003. Effect of temperature on lipids, proteins, and carbohydrates levels during development from egg extrusion to juvenile stage of *Cherax quadricarinatus* (Decapoda: Parastacidae). *Comp. Biochem. Physiol.* 138, 147–154.
- Grasshoff, K., 1983. *Methods of Seawater Analysis*. Verlag chemie, NY, pp. 317.
- Guerreiro, I., Peres, H., Castro-Cunha, M., Oliva-Teles, A., 2012. Effect of temperature and dietary protein/lipid ratio on growth performance and nutrient utilization of juvenile Senegalese sole (*Solea senegalensis*). *Aquac. Nutr.* 18, 98–106.
- Guinea, J., Fernandez, F., 1997. Effect of feeding frequency, relative meal size and temperature on energy metabolism in *Sparus aurata*. *Aquaculture* 148, 125–142.
- Hasan, M.R., Macintosh, D.J., 1991. Effect of environmental temperature and feeding rate on the growth, food utilization and body composition of common carp (*Cyprinus carpio* L.) fry. In: Kaushik, S.J., Luquet, P. (Eds.), *Fish Nutrition in Practice*. Les Colloques, Paris, pp. 767–778.
- Huang, J.F., Xu, Q.Y., Chang, Y.M., 2016. Effects of temperature and dietary protein on the growth performance and *IGF-I* mRNA expression of juvenile mirror carp (*Cyprinus carpio*). *Aquac. Nutr.* 22, 283–292.
- Iannucci, F.M., 2015. *Deeper Effects of Climate Change: The Influence of Temperature and Diet Quality on Fish Metabolism*. UVM Honors College Senior Theses, pp. 89.
- IPCC, 2014. *Climate change 2014: impacts, adaptation, and vulnerability*. Part A: global and sectoral aspects. In: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R., White, L.L. (Eds.), *Contribution of Working Group II to the Fifth. Press*, Cambridge, United Kingdom and New York, NY, USA, pp. 11–32.
- Jadhav, M., Bawane, V., Gulave, A., 2012. Size dependent variation in the rate of oxygen consumption, ammonia and O:N ratio of freshwater bivalve, *Lamellidens marginalis* from Godavari river during monsoon (M.S) India. *Trends Fish. Res.* 1 (2), 22–26.
- Jobling, M., 1994. *Fish Bioenergetics*. Chapman & Hall, London (309 pp).
- Kao, Y.-C., Madenjian, C.P., Bunnell, D.B., Lofgren, B.M., Perroud, M., 2015. Temperature effects induced by climate change on the growth and consumption by salmonines in Lakes Michigan and Huron. *Environ. Biol. Fishes* 98, 1089–1104.
- Kierkegaard, A., Balk, L., Tjarnlund, U., De Wit, C.A., Jansson, B., 1999. Dietary uptake and biological effects of decabromodiphenyl ether in rainbow trout (*Oncorhynchus mykiss*). *Environ. Sci. Technol.* 33, 1612–1617.
- Kreiss, C.M., Michael, K., Lucassen, M., Jutfelt, F., Motyka, R., Dupont, S., Pörtner, H.-O., 2015. Ocean warming and acidification modulate energy budget and gill ion regulatory mechanisms in Atlantic cod (*Gadus morhua*). *J. Comp. Physiol. B* 185, 767–781.
- Kuo, Y.-M., Sepúlveda, M.S., Sutton, T.M., Ochoa-Acuña, H.G., Muir, A.M., Miller, B., Hua, I., 2010. Bioaccumulation and biotransformation of decabromodiphenyl ether and effects on daily growth in juvenile lake whitefish (*Coregonus clupeaformis*). *Ecotoxicology* 19, 751–760.
- Lemos, D., Netto, B., Germano, A., 2006. Energy budget of juvenile fat snook *Centropomus parallelus* fed live food. *Comp. Biochem. Physiol. Part A: Mol. Integr. Physiol.* 144, 33–40.
- Leung, K.M.Y., Chu, J.C.W., Wu, R.S.S., 1999. Effects of body weight, water temperature an ration size on ammonia excretion by the areolated group (*Epinephelus areolatus*) and mangrove snapper (*Lutjanus argentimaculatus*). *Aquaculture* 170, 215–227.
- Mace, A.M., Ansell, A.D., 1982. Respiration and nitrogen excretion of *Polinices alderi* (Forbes) and *Polinices catens* (da Costa) (Gastropoda: Naticidae). *J. Exp. Mar. Biol. Ecol.* 60 (3), 275–292.
- Marques, A., Nunes, M.L., Moore, S.K., Strom, M.S., 2010. Climate change and seafood safety: human health implications. *Food Res. Int.* 43 (7), 1766–1779.
- Maulvault, A.L., Barbosa, V., Alves, R., Custódio, A., Anacleto, P., Repolho, T., Ferreira, P.P., Rosa, R., Marques, A., Diniz, M., 2017. Ecophysiological responses of juvenile seabass (*Dicentrarchus labrax*) exposed to increased temperature and dietary methylmercury. *Sci. Total Environ.* 586, 551–558.
- Maulvault, A.L., Custódio, A., Anacleto, P., Repolho, T., Pousão, P., Nunes, M.L., Diniz, M., Rosa, R., Marques, A., 2016. Bioaccumulation and elimination of mercury in juvenile seabass (*Dicentrarchus labrax*) in a warmer environment. *Environ. Res.* 149, 77–85.
- Morato, T., Afonso, P., Lourinho, P., Nash, R.D.M., Santos, R.S., 2003. Reproductive biology and recruitment of the white seabream in the Azores. *J. Fish Biol.* 63, 59–72.
- Noyes, P.D., Hinton, D.E., Stapleton, H.M., 2011. Accumulation and debromination of decabromodiphenyl ether (BDE-209) in juvenile fathead minnows (*Pimephales promelas*) induces thyroid disruption and liver alterations. *Toxicol. Sci.* 122 (2), 265–274.
- Noyes, P.D., Stapleton, H.M., 2014. PBDE flame retardants: Toxicokinetics and thyroid hormone endocrine disruption in fish. *Endocr. Disruptors* 2 (1), e29430.
- Pang, X., Yuan, X.-Z., Cao, Z.-D., Zhang, Y.-G., Fu, S.-J., 2015. The effect of temperature on repeat swimming performance in juvenile qingbo (*Spirinbarbus sinensis*). *Fish Physiol. Biochem.* 41, 19–29.
- Peres, M.H., Oliva-Teles, A., 1999. Effect of dietary lipid level on growth performance and feed utilization by European sea bass juveniles (*Dicentrarchus labrax*). *Aquaculture* 179, 325–334.
- Pörtner, H.O., Knust, R., 2007. Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* 315, 95–97.
- Pörtner, H.O., Peck, M.A., 2010. Climate change effects on fishes and fisheries: towards a cause-and-effect understanding. *J. Fish Biol.* 77, 1745–1779.
- Requena, A., Fernández-Borrás, J., Planas, J., 1997. The effects of a temperature rise on oxygen consumption and energy budget in gilthead sea bream. *Aquac. Int.* 5, 415–426.
- Ricker, W.E., 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Can.* 191, 1–382.
- Rocha, A.J.S., Gomes, V., Phan, V.N., Passos, M.J.A.C.R., Furia, R.R., 2005. Metabolic demand and growth of juveniles of *Centropomus parallelus* as function of salinity. *J. Exp. Mar. Biol. Ecol.* 316, 157–165.
- Rosa, R., Pimentel, M.S., Boavida-Portugal, J., Teixeira, T., Trübenbach, K., Diniz, M.S., 2012b. Ocean warming enhances malformations, premature hatching, metabolic suppression and oxidative stress in the early life stages of a keystone invertebrate. *PLoS One* 7 (6), e38282.
- Rosa, R., Marques, A., Nunes, M.L., 2012a. Impact of climate change in Mediterranean aquaculture. *Rev. Aquac.* 4, 163–177.
- Rosa, R., Seibel, B.A., 2008. Synergistic effects of climate-related variables suggest future physiological impairment in a top oceanic predator. *Proc. Natl. Acad. Sci. USA* 105, 20776–20780.
- Rosa, R., Seibel, B.A., 2010. Voyage of the argonauts in the pelagic realm: physiological and behavioral ecology of the rare paper nautilus *Argonauta nouryi*. *ICES J. Mar. Sci.* 67 (7), 1494–1500.
- Rosa, R., Lopes, A.R., Pimentel, M., Faleiro, F., Baptista, M., Trübenbach, K., Narciso, L., Dionísio, G., Pegado, M.R., Repolho, T., Calado, R., Diniz, M., 2014. Ocean cleaning stations under a changing climate: biological responses of tropical and temperate fish-cleaner shrimp to global warming. *Glob. Change Biol.* 20 (10), 3068–3079.
- Sampaio, S., Lopes, A.R., Francisco, S., Paula, J.R., Pimentel, M., Maulvault, A.L., Repolho, T., Grilo, T.F., Pousão-Ferreira, P., Marques, A., Rosa, R., 2018. Ocean acidification dampens physiological stress response to warming and contamination in a commercially-important fish (*Argyrosomus regius*). *Sci. Total Environ.* 618, 388–398.
- Sandersfeld, T., Davison, W., Lamare, M.D., Knust, R., Richter, C., 2015. Elevated temperature causes metabolic trade-offs at the whole-organism level in the Antarctic fish, *Trematomus bernacchii*. *J. Exp. Biol.* 218, 2373–2381.

- Sha, J., Wang, Y., Lv, J., Wang, H., Chen, H., Qi, L., Tang, X., 2015. Effects of two polybrominated diphenyl ethers (BDE-47, BDE-209) on the swimming behavior, population growth and reproduction of the rotifer *Brachionus plicatilis*. *J. Environ. Sci.* 28, 54–63.
- Smith, L.S., 1989. Digestive functions in teleost fishes. In: Halver, J.E. (Ed.), *Fish Nutrition*. Academic Press, New York, USA, pp. 331–421.
- Stapleton, H.M., Brazil, B., Holbrook, R.D., Mitchelmore, C.L., Benedict, R., Konstantinov, A., Potter, D., 2006. In vivo and in vitro debromination of decabromodiphenyl ether (BDE 209) by juvenile rainbow trout and common carp. *Environ. Sci. Technol.* 40 (15), 4653–4658.
- Sun, L., Chen, H., Huang, L., 2006. Effect of temperature on growth and energy budget of juvenile cobia (*Rachycentron canadum*). *Aquaculture* 261, 872–878.
- Sun, L., Chen, H., 2009. Effects of ration and temperature on growth, fecal production, nitrogenous excretion and energy budget of juvenile cobia (*Rachycentron canadum*). *Aquaculture* 292, 197–206.
- Tang, Q., Sun, Y., Zhang, B., 2003. Bioenergetics models for seven species of marine fish. *J. Fish. Chin.* 27 (5), 443–449 (in Chinese with English abstract).
- Tidwell, J.H., Coyle, S.D., Bright, L.A., Amum, A.V., Yasharian, D., 2003. Effect of water temperature on growth, survival, and biochemical composition of largemouth bass *Micropterus salmoides*. *J. World Aquac. Soc.* 34 (2), 175–183.
- Van Ham, E.H., Berntssen, M.H.G., Imsland, A.K., Parpoura, A.C., Bonga, S.E.W., Stefansson, S.O., 2003. The influence of temperature and ration on growth, feed conversion, body composition and nutrient retention of juvenile turbot (*Scophthalmus maximus*). *Aquaculture* 217, 547–558.
- Windisch, H.S., Kathöver, R., Pörtner, H.O., Frickenhaus, S., Lucassen, M., 2011. Thermal acclimation in Antarctic fish: transcriptomic profiling of metabolic pathways. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301 (5), R1453–R1466.
- Xie, S., Cui, Y., Yang, Y., Liu, J., 1997. Energy budget of Nile tilapia (*Oreochromis niloticus*) in relation to ration size. *Aquaculture* 154, 57–68.
- Xie, X.J., Sun, R., 1992. The bioenergetics of the southern catfish (*Silurus meridionalis* Chen): growth rate as a function of ration level, body weight, and temperature. *J. Fish Biol.* 40 (5), 719–730.
- Xie, X.J., Sun, R., 1993. Pattern of energy allocation in the southern catfish (*Silurus meridionalis* Chen). *J. Fish Biol.* 42, 197–207.
- Xie, S., Zheng, K., Chen, J., Zhang, Z., Zhu, X., Yang, Y., 2011. Effect of water temperature on energy budget of Nile tilapia, *Oreochromis niloticus*. *Aquac. Nutr.* 17, e683–e690.
- Yang, Y.O., Cui, Y.B., Xiong, B.X., Yang, Y.X., Xie, S.Q., 2003. Comparative studies on nitrogen budget and energy budget of Jian carp (*Cyprinus carpio* var. jian) and gibel carp (*Carassius auratus gibelio*) fed diets with different qualities. *Acta Hydrobiol. Sin.* 27 (6), 572–579 (in Chinese with English abstract).
- Ye, L., Jiang, S.G., Zhu, X.M., Yang, Q.B., Wen, W.G., Wu, K.C., 2009. Effects of salinity on growth and energy budget of juvenile *Penaeus monodon*. *Aquaculture* 290, 140–144.
- Yurista, P.M., 1999. Temperature-dependent energy budget of an Arctic cladoceran, *Daphnia middendorffiana*. *Freshw. Biol.* 42, 21–34.
- Zhang, X., Tang, X., Zhou, B., Wang, Y., 2013. Effects of decabromodiphenyl ether (BDE-209) on inter-specific competition between two species of marine bloom-forming microalgae. *PLoS One* 8 (3), e56084.
- Zheng, Z., Jin, C., Li, M., Bai, P., Dong, S., 2008. Effects of temperature and salinity on oxygen consumption and ammonia excretion of juvenile miuy croaker, *Miichthys miuy* (Basilewsky). *Aquac. Int.* 16, 581–589.